



Cost-Effectiveness of a Diabetes Self-Management Education and Support Intervention Led by Community Health Workers and Peer Leaders: Projections From the Racial and Ethnic Approaches to Community Health Detroit Trial

Diabetes Care 2021;44:1108–1115 | <https://doi.org/10.2337/dc20-0307>

Wen Ye,¹ Shihchen Kuo,² Edith C. Kieffer,³
Gretchen Piatt,⁴ Brandy Sinco,³
Gloria Palmisano,⁵ Michael S. Spencer,⁶
and William H. Herman²

OBJECTIVE

To simulate the long-term cost-effectiveness of a peer leader (PL)-led diabetes self-management support (DSMS) program following a structured community health worker (CHW)-led diabetes self-management education (DSME) program in reducing risks of complications in people with type 2 diabetes (T2D).

RESEARCH DESIGN AND METHODS

The trial randomized 222 Latino adults with T2D to 1) enhanced usual care (EUC); 2) a CHW-led, 6-month DSME program and 6 months of CHW-delivered monthly telephone outreach (CHW-only); or 3) a CHW-led, 6-month DSME program and 12 months of PL-delivered weekly group sessions with telephone outreach to those unable to attend (CHW + PL). Empirical data from the trial and the validated Michigan Model for Diabetes were used to estimate cost and health outcomes over a 20-year time horizon from a health care sector perspective, discounting both costs and benefits at 3% annually. The primary outcome measure was the incremental cost-effectiveness ratio (ICER).

RESULTS

Over 20 years, the CHW + PL intervention had an ICER of \$28,800 and \$5,900 per quality-adjusted life-year (QALY) gained compared with the EUC and CHW-only interventions, respectively. The CHW-only intervention had an ICER of \$430,600 per QALY gained compared with the EUC intervention. In sensitivity analyses, the results comparing the CHW + PL with EUC and CHW-only interventions were robust to changes in intervention effects and costs.

CONCLUSIONS

The CHW + PL-led DSME/DSMS intervention improved health and provided good value compared with the EUC intervention. The 6-month CHW-led DSME intervention without further postintervention CHW support was not cost-effective in Latino adults with T2D.

¹Department of Biostatistics, University of Michigan School of Public Health, Ann Arbor, MI

²Department of Internal Medicine, University of Michigan Medical School, Ann Arbor, MI

³University of Michigan School of Social Work, Ann Arbor, MI

⁴Department of Learning Health Sciences, University of Michigan Medical School, Ann Arbor, MI

⁵Community Health and Social Service Center, Inc., Detroit, MI

⁶University of Washington School of Social Work, Seattle, WA

Corresponding author: Wen Ye, wye@umich.edu

Received 13 February 2020 and accepted 22 February 2021

This article contains supplementary material online at <https://doi.org/10.2337/figshare.14098814>.

W.Y. and S.K. have equal first authorship.

© 2021 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <https://www.diabetesjournals.org/content/license>.

Approximately 30 million people in the U.S. have diabetes, and 1.5 million are diagnosed with diabetes each year (1). The total cost of diagnosed diabetes in the U.S. reached \$327 billion in 2017, accounting for 24% of all health care expenditures (2). Evidence has demonstrated the short-term (e.g., 6 month) effectiveness of diabetes self-management education (DSME) in improving clinical outcomes and quality of life (3–6). DSME is associated with increased use of primary and preventive services and lower use of inpatient hospital services and is cost-effective (7). The American Diabetes Association recommends that all individuals with type 2 diabetes (T2D) receive DSME. However, given the number of people with diabetes, the limited number of diabetes educators, and inequities in access to high-quality care, the current health system cannot support long-term DSME (8).

Efforts have shifted to community resources to meet these challenges (9). Increasingly, community health workers (CHWs) have been used to provide structured DSME, especially in low-income and racial/ethnic minority communities and populations (10) where effective, culturally tailored approaches to diabetes self-management are needed (11,12). CHWs are trusted members of their communities and may relate to and empathize with their clients on a level that a credentialed health care professional may not (13,14).

Diabetes self-management support (DSMS) is defined as “activities that assist the individual with diabetes to implement and maintain the on-going behaviors needed to manage their illness.” (5) The type of support provided can be behavioral, educational, psychosocial, and/or clinical in nature. Although CHWs have been shown to improve short-term self-monitoring, self-care, lifestyle change, and blood glucose control (5,6,13,15,16), it has been challenging to maintain the gains over the long term. Most patients require DSMS. One promising approach is to offer less intensive support services using peer leaders (PLs) after intensive CHW interventions (6,13,16–20). PLs share important characteristics with participants, including having diabetes. They are often volunteers or receive small stipends to reimburse any expenses (17,18,21). Some PL-led

interventions have resulted in improvements in HbA_{1c}, self-empowerment scores, self-care indicators (16), patient activation (17), and diabetes-related distress (17). The most effective diabetes PL models offer support following structured DSME (6,10,18,20), are delivered through multiple modes of interaction (10), and are implemented in community settings.

The Racial and Ethnic Approaches to Community Health (REACH) Detroit Partnership, in collaboration with a federally qualified health center (Community Health and Social Services Center [CHASS]) adapted a DSME curriculum from the Diabetes Prevention Program using a community-based participatory approach (11,12), implemented it using trained CHWs, and tested it over several phases (5,6,22). Between October 2009 and February 2013, CHASS and the University of Michigan used this curriculum (Journey to Health/El Camino a la Salud) to conduct a cluster randomized clinical trial of a CHW-led DSME intervention to improve diabetes self-management, glycemic control, and other clinical and psychosocial outcomes among Latino adults with T2D receiving health care at CHASS (6). The trial enrolled 222 individuals with T2D who were first randomized to the CHW-led, 6-month DSME program ($n = 149$) or enhanced usual care (EUC) ($n = 73$). The CHW intervention included 1) 11 Journey to Health/El Camino a la Salud DSME classes, 2) a CHW-accompanied clinic visit to the participant and his or her primary care provider, 3) support from the CHW for goal setting, and 4) home visits and biweekly phone calls. The CHWs were Latinas from the southwest Detroit community who were fluent in Spanish and had received core competency CHW training, diabetes education training, and behavior modification strategies (6). The EUC group attended a 2-h class conducted by a research assistant that covered the interpretation of clinical and anthropometric information. Both groups received usual care at CHASS. At the end of the 6-month DSME program, the CHW-led group was further randomized to receive 1) brief monthly check-in phone calls from the CHW between 7 and 12 months and no CHW contact from 13 to 18 months (CHW-only) ($n = 89$) or 2) 12 months of weekly in-person group sessions of DSMS delivered by

PLs (CHW + PL) ($n = 60$). All groups, including the EUC group, continued to receive usual care. The PLs were prior Journey to Health/El Camino a la Salud participants who completed PL training, received monthly booster sessions, and were supervised by the CHWs. This PL intervention component was added to the original design of the trial as a pilot study. The trial showed that participants in the CHW intervention at the 6-month follow-up had significant reductions in HbA_{1c} and diabetes distress compared with EUC. CHW + PL participants maintained HbA_{1c} improvements at 12 and 18 months, and CHW-only participants maintained improvements in diabetes distress at 12 and 18 months. Details of the study design, interventions, and results have been published elsewhere (6).

To date, for the management of T2D, there is no published research evaluating the cost-effectiveness of CHW + PL DSME/DSMS interventions, and there are limited studies that assessed the cost-effectiveness of CHW-only interventions (23). We sought to extend the trial results using a computer simulation model to estimate the longer term costs and health outcomes and to assess the long-term cost-effectiveness of the interventions, especially the cost-effectiveness of adding PL DSMS after a short-term intensive CHW-led DSME intervention.

RESEARCH DESIGN AND METHODS

Simulation Model

We used a validated microsimulation model for T2D, the Michigan Model for Diabetes (MMD) version 2.0 (24–27), to simulate the long-term cost-effectiveness of the CHW + PL intervention compared with EUC and the CHW-only intervention. Disease progression in the MMD is based on six discrete-time discrete-event submodels that simulate diabetes-related complications (retinopathy, nephropathy, neuropathy), major comorbidities (coronary heart disease, cerebrovascular disease), and death. Transition probabilities in these submodels are functions of individual characteristics, risk factor levels, and current disease and treatment states. The model also estimates the costs of diabetes and the health-related quality of life associated with the health states. Details

of the MMD have been published elsewhere (24–27).

Simulation Population Characteristics and Modeled Health Outcomes

We incorporated information about the characteristics of the REACH Detroit trial participants and the effectiveness of the trial interventions into the MMD. The published analyses revealed that at 6 months after randomization, HbA_{1c} decreased significantly more in participants who received the CHW intervention compared with those who received EUC (6). From 6 to 12 and 12 to 18 months, improvements in HbA_{1c} were sustained for participants randomized to the CHW + PL group but not for those randomized to the EUC or the CHW-only groups (6). In addition, significant decreases in systolic blood pressure (SBP), diastolic blood pressure (DBP), and LDL cholesterol (LDLc) were observed in the CHW + PL group at 18 months compared with baseline. The time horizon for the simulation began at the end of the REACH Detroit trial (i.e., 18 months following randomization). For risk factors with significant within- or between-group differences (i.e., HbA_{1c}, SBP, DBP, and LDLc), group-specific summary statistics at 18 months were used as the initial simulation population characteristics. For all other variables (e.g., BMI, HDL cholesterol [HDLc], health utility scores), summary statistics that are based on the pooled population at 18 months were used for the simulation (Supplementary Table 1).

Continuous measures at 18 months following randomization were simulated using a Gaussian distribution with the given mean (median) and SD but truncated at ± 3 SDs. Height and weight were simulated with a correlation of 0.51. SBP and DBP were simulated with a correlation of 0.82. Highly skewed distributions, such as HbA_{1c} and triglyceride levels, were simulated using log-normal distributions.

To evaluate the long-term effectiveness and cost-effectiveness of the interventions, we estimated the incidence of clinical outcomes, life expectancy, and quality-adjusted life expectancy (QALE). The seven clinical outcomes included three-point major adverse cardiovascular events (MACE) (including nonfatal myocardial infarction [MI], nonfatal

stroke, or cardiovascular death) and defined as the primary outcome, fatal or nonfatal MI, fatal or nonfatal stroke, coronary revascularization procedures, hospitalization for heart failure, death as a result of cardiovascular causes, and all-cause mortality.

Intervention and Outcome Costs

We considered both the costs of implementing the interventions (intervention costs) and the medical costs of T2D and its complications (outcome costs) in the cost-effectiveness analyses. To estimate intervention costs, we included the costs of 1) the 2-h class for the EUC group, 2) identifying and training PLs, and 3) implementing and maintaining CHW-only and CHW + PL interventions. These costs were summed to derive the total intervention costs. The per-participant intervention costs were \$1,599, \$820, and \$234 for the CHW + PL, CHW-only, and EUC groups, respectively, over 18 months (Supplementary Table 2).

Outcome costs refer to the medical costs of T2D and its complications (Supplementary Table 3). We did not consider the medical costs of care incurred or averted by the interventions within the 18 months of the trial because the observed utilization of medical services (hospital, emergency department, and outpatient visits) and the use of prescription medications for hyperglycemia, hypertension, and dyslipidemia were not significantly different among the CHW + PL, CHW-only, and EUC groups. Outcome costs were estimated from the MMD on the basis of the published literature (28). Costs were assessed according to sex, BMI, antihyperglycemic therapy, and diabetes-related complications. The costs of complications were estimated as the cost incurred during the first year that a complication occurred (event cost) and the cost in each subsequent year after the complication occurred (ongoing cost). We also considered the costs of death.

Following the recommendations of the Second Panel on Cost-Effectiveness in Health and Medicine (29), we performed cost-effectiveness analyses from the health care sector perspective that considered medical costs (the intervention and outcome costs) borne by third-party payers. The impact inventory is

provided in Supplementary Table 4. All costs were expressed in 2018 U.S. dollars.

Health Utilities

Health utility scores are a measure of health-related quality of life in which perfect health is assigned a value of 1.0 and death a value of 0.0. In economic analyses, the health utility score for each health state is multiplied by the time a subject spends in that health state. These are then summed to calculate the QALE, which is expressed as quality-adjusted life-years (QALYs) accrued over a specified period of time. The MMD incorporates a health utility module to calculate yearly and cumulative QALYs on the basis of subjects' demographic, diabetes treatment, complication, and comorbidity status (27,30).

Base-Case Analysis and Sensitivity Analyses

In the base-case analysis, we assumed that no additional interventions were implemented after 18 months and that the effects of the interventions observed at 18 months diminished over the simulation time horizon. Under this scenario, intervention costs only included the costs incurred over the 18-month trial period. We projected trajectories of HbA_{1c}, blood pressure, and lipid ratio (total cholesterol/HDLc) after 18 months according to the equations derived from the UK Prospective Diabetes Study Outcomes Model (UKPDS-OM) (31), which demonstrated gradual convergence of each risk factor as the treatment effect/benefit wore off.

In all analyses, we used a 20-year simulation time horizon. We also ran the MMD simulation for 10 years and 30 years to assess the shorter- and longer-term cost-effectiveness of the interventions, respectively. We minimized the first-order uncertainty (stochastic uncertainty) by performing Monte Carlo simulations with a large enough number of replications (3,000,000) for each of the three intervention groups so that the mean simulated event rates were stable (SEs ~ 0.01 – 0.03). We assessed the cost-effectiveness of the CHW + PL intervention relative to the EUC and CHW-only interventions using incremental cost-effectiveness ratios (ICERs) calculated as incremental costs divided by

incremental QALYs. We also assessed the cost-effectiveness of the CHW-only intervention relative to EUC. We discounted both costs and QALYs at 3% per year.

Second-order uncertainty (parameter uncertainty) was assessed through sensitivity analyses. In the first two sensitivity analysis scenarios, we assumed that the intervention effects observed at 18 months persisted over the simulation time horizon and that the levels of risk factors (i.e., HbA_{1c}, blood pressure, lipids) did not change over time. In the first scenario, we assumed that no additional resources were needed to maintain the level of these risk factors and that the intervention costs included only the costs incurred over the 18-month trial period. This is an optimistic assumption. In the second scenario, we assumed that resources were needed to maintain the intervention benefit after 18 months. Therefore, we included ongoing intervention costs and assumed that 2 h per participant per year were required for the PL or CHW over the simulation time horizon. Under the assumption that effects of the interventions observed at 18 months diminished over the simulation time horizon (same as in the base-case analysis), we also performed one-way sensitivity analyses to assess the impact of cost and treatment effect parameters on the 20-year cost-effectiveness of the CHW + PL versus the EUC interventions (Supplementary Table 5). For the cost parameter, we assumed a 50% reduction or a 50% increase in intervention costs for the CHW + PL intervention. For the treatment effect parameters, the mean initial level of HbA_{1c} in the CHW + PL group was increased or decreased by 1 SE of the estimated difference in change of HbA_{1c} from randomization to 18 months postrandomization between the CHW + PL and EUC groups. The mean levels of SBP and LDLc were adjusted in the same manner.

All simulations were performed using Indirect Estimation and Simulation Tools software (32). Results from the simulation of the three intervention groups were evaluated using SAS statistical software (SAS Institute, Cary, NC).

There are no absolute criteria for cost-effectiveness in the U.S., and the long-cited benchmark of \$50,000 per QALY gained for an intervention to be deemed cost-effective is largely

unsupported (33). Nevertheless, in general, interventions costing <\$20,000 per QALY gained may be considered as having strong evidence for adoption, interventions costing \$20,000–\$100,000 per QALY gained have moderate evidence for adoption, and those costing >\$100,000 per QALY gained have weaker evidence for adoption (34). It has been suggested that a threshold of \$100,000 or \$150,000 per QALY gained may better reflect the U.S. health care environment (35) and represent good value for money.

RESULTS

Supplementary Table 1 shows the characteristics of the three study groups at the end of the 18-month trial and the characteristics of the three simulated groups at the initiation of the model simulation. The simulated population of adults with T2D reflected the pooled sample means from the three study groups. The simulated Latino population had a mean age of 50.4 years, 39% were male, mean T2D duration was 7.6 years, and mean BMI was 32.4 kg/m². At the end of the trial, the CHW + PL group had the lowest HbA_{1c}, SBP, DBP, and LDLc; the CHW-only group had the highest SBP and DBP; and the EUC group had the highest HbA_{1c} and LDLc. Figure 1 shows the gradual convergence of risk factors for diabetes progression over the simulation period in the base-case scenario. The magnitude of the difference in risk factors among the three groups become smaller over time.

Health Outcomes

Table 1 summarizes the simulated 20-year cumulative incidence of diabetes complications by intervention group. In the base-case scenario with diminishing intervention benefit overtime, clinical outcomes were generally best for the CHW + PL intervention group and worse for the CHW-only and EUC intervention groups. For example, the cumulative incidence of three-point MACE was lowest in the CHW + PL group (30.1%) and higher in the CHW-only (30.5%) and EUC groups (30.6%). In the scenario in which we assumed persistent intervention benefits, clinical outcomes were best for the CHW + PL




Figure 1—Simulated risk factor trajectories for the scenario assuming the diminishing intervention benefit after the end of the REACH Detroit trial and over the simulation period.

intervention group and worse for the CHW-only or EUC intervention groups. For example, the cumulative incidence of three-point MACE was lowest in the CHW + PL group (23.6%) followed by the CHW (26.1%) and EUC groups (26.9%). The cumulative incidence of stroke was lowest in the CHW + PL group (5.1%) followed by the EUC group (6.0%) and CHW-only groups (6.1%).

Cost-Effectiveness

Table 2 summarizes the simulated outcomes for the base-case analysis. Over a 20-year period, the CHW + PL

Table 1—Cumulative incidence of complications when intervention effects diminish or persist over 20 years by intervention group

Complication	Scenario with diminishing intervention effects ¹ , %			Scenario with persistent intervention effects ² , %		
	EUC	CHW only	CHW + PL	EUC	CHW only	CHW + PL
MI	14.5	14.4	14.1	12.6	12.0	10.7
Stroke	7.3	7.3	7.1	6.0	6.1	5.1
Cardiovascular death	21.0	20.9	20.6	18.6	18.0	16.1
Three-point MACE	30.6	30.5	30.1	26.9	26.1	23.6
Revascularization procedure	24.9	24.7	24.3	22.0	21.1	19.3
Congestive heart failure	18.1	18.2	18.1	17.3	17.2	16.3
All-cause death	34.3	34.3	34.0	32.2	31.6	30.0

¹We used the UKPDS-OM risk equations to project gradual convergence trajectories of the postintervention levels of all risk factors for diabetes complications (i.e., HbA_{1c}, SBP, and lipid ratio [total cholesterol divided by HDLc]) as the treatment effect wears off over a period of several years. ²We assumed that the postintervention levels of all risk factors for diabetes complications (i.e., HbA_{1c}, SBP, and lipid ratio [total cholesterol divided by HDLc]) did not change over the simulation time horizon.

intervention was associated with the longest life expectancy (17.06 years) and QALE (7.07 QALYs). The EUC intervention resulted in the lowest total costs over 20 years (\$127,639). Compared with the EUC group, the CHW + PL group cost \$796 more and gained 0.028 more QALYs, leading to an ICER of ~\$28,800 per QALY gained. Compared with the CHW-only group, the CHW + PL group cost ~\$5,900 per QALY gained. Compared with the EUC group, the CHW-only group cost \$641 more

and gained 0.002 QALYs, resulting in an ICER of ~\$430,600 per QALY gained. Over the 10-year and 30-year time horizons, compared with the EUC intervention, the CHW + PL group cost ~\$44,200 and \$28,200 per QALY gained, respectively, and the CHW-only group cost ~\$4,443,300 and \$402,600 per QALY gained, respectively.

Table 3 summarizes the results of the two sensitivity analyses. When we assumed that the intervention benefit observed at 18 months persisted over 20

years without further intervention, the CHW + PL group dominated both the EUC and the CHW-only groups. That is, the CHW + PL group cost \$1,933 less and gained 0.092 more QALYs than the EUC group and cost \$1,798 less and gained 0.071 more QALYs than the CHW-only group. Compared with the EUC group, the CHW-only group was also cost saving over 20 years. Compared with the EUC group, the CHW-only group produced an ICER of ~\$5,100 per QALY gained over 10 years

Table 2—Base-case analysis modeling cost-effectiveness of interventions for 18 months with diminishing intervention benefits after 18 months

	EUC	CHW only	CHW + PL	CHW + PL vs. CHW only	CHW only vs. EUC	CHW + PL vs. EUC
Intervention cost ¹ (\$)	234	820	1,599	779	586	1,365
Outcome cost (\$)						
10 years	71,934	71,899	71,226	−674	−35	−708
20 years	127,405	127,460	126,836	−624	55	−569
30 years	163,073	163,164	162,644	−520	92	−428
Total cost (\$)						
10 years	72,168	72,719	72,825	105	551	657
20 years	127,639	128,280	128,435	155	641	796
30 years	163,307	163,984	164,243	259	678	937
Life expectancy (years)						
10 years	9.35	9.36	9.38	0.019	0.0040	0.0236
20 years	17.01	17.02	17.06	0.0478	0.0091	0.0569
30 years	22.65	22.66	22.72	0.0692	0.0096	0.0788
QALE (QALYs)						
10 years	4.46	4.46	4.48	0.0147	0.0001	0.0149
20 years	7.05	7.05	7.07	0.0262	0.0015	0.0276
30 years	8.40	8.41	8.44	0.0316	0.0017	0.0333
Cost per QALY gained (\$)						
10 years				7,168	4,443,300	44,172
20 years				5,918	430,600	28,796
30 years				8,202	402,594	28,164

¹Intervention cost was the per-participant cost over the 18-month intervention period of the REACH Detroit trial.

Table 3—Sensitivity analyses modeling cost-effectiveness of interventions

Sensitivity analysis scenario	EUC	CHW only	CHW + PL	CHW + PL vs. CHW only	CHW only vs. EUC	CHW + PL vs. EUC
18-month intervention with persistent intervention benefits						
Total cost ¹ (\$)						
10 years	70,851	70,878	69,985	−893	27	−866
20 years	124,401	124,265	122,467	−1,798	−135	−1,933
30 years	159,312	159,367	157,491	−1,876	54	−1,822
QALE (QALYs)						
10 years	4.48	4.49	4.51	0.0234	0.0051	0.0286
20 years	7.14	7.16	7.23	0.0714	0.0206	0.0919
30 years	8.58	8.61	8.73	0.1183	0.0348	0.1532
Cost per QALY gained (\$)						
10 years				Cost saving	5,148	Cost saving
20 years				Cost saving	Cost saving	Cost saving
30 years				Cost saving	1,570	Cost saving
Ongoing intervention with persistent intervention benefits						
Total cost ² (\$)						
10 years	70,851	71,252	70,545	−706	400	−306
20 years	124,401	124,867	123,371	−1,497	467	−1,029
30 years	159,312	160,109	158,604	−1,505	796	−709
QALE (QALYs)						
10 years	4.48	4.49	4.51	0.0234	0.0051	0.0286
20 years	7.14	7.16	7.23	0.0714	0.0206	0.0919
30 years	8.58	8.61	8.73	0.1183	0.0348	0.1532
Cost per QALY gained (\$)						
10 years				Cost saving	78,148	Cost saving
20 years				Cost saving	22,719	Cost saving
30 years				Cost saving	22,862	Cost saving

¹Total cost was a combination of the intervention cost and outcome cost. The intervention cost was \$234, \$820, and \$1,599 per participant for the EUC, CHW-only, and CHW + PL interventions, respectively, over the 18-month intervention period of the REACH Detroit trial. The outcome cost derived from the direct medical costs of diabetes and its complications was estimated by the MMD. ²Total cost was a combination of the intervention cost and outcome cost. The intervention cost was \$234, \$820, and \$1,599 per participant for the EUC, CHW-only, and CHW + PL interventions, respectively, over the 18-month intervention period of the REACH Detroit trial. The ongoing intervention cost per participant per year was \$43.80 and \$65.70 for the CHW-only and CHW + PL interventions over the simulation period, respectively, assuming that a total of 2 h per participant per year was required by the CHW or PL to make contact with the participants. The ongoing cost would support the ongoing CHW-only and CHW + PL interventions to maintain the trial-observed intervention effects over the simulation time horizon. The outcome cost derived from the direct medical costs of diabetes and its complications was estimated by the MMD.

and \$1,600 per QALY gained over 30 years. Over the 10-year and 30-year time horizons, the CHW + PL group still dominated the other two groups.

In the second sensitivity analysis, we assumed that limited ongoing interventions were needed after 18 months to maintain the benefits observed at 18 months (Table 3). Under this assumption, over 20 years, the CHW + PL dominated both the EUC and the CHW-only groups. The CHW-only group compared with the EUC group yielded an ICER of ~\$22,700 per QALY gained. Over the 10-year and 30-year time horizons, the CHW + PL group still dominated the EUC and CHW-only groups. The CHW-only group cost ~\$78,100 and ~\$22,900 per QALY gained, respectively, compared with the EUC group.

Supplementary Table 5 shows the effects of plausible changes in the intervention costs and treatment effects of the CHW + PL intervention on cost-effectiveness under the assumption that effects of the interventions observed at 18 months diminished over the simulation time horizon. Compared with the EUC intervention, a 50% reduction in the intervention costs would make the CHW + PL intervention cost saving. A 50% increase in the intervention costs would increase the ICER of the CHW + PL intervention to ~\$57,700 per QALY gained. When we assumed a smaller treatment effect on one of the three key risk factors, the ICER of CHW + PL versus EUC increased, ranging from \$37,600 to \$46,092 per QALY gained. When assuming a larger

treatment effect on one of the three key risk factors, the ICER decreased, ranging from \$15,000 to \$18,400 per QALY gained.

CONCLUSIONS

In this study, we showed that a CHW + PL-led DSME/DSMS intervention in Latino adults with T2D and poor glycemic control in a low-income urban setting was cost-effective compared with EUC. In the base-case scenario, compared with the EUC strategy, the CHW + PL strategy cost ~\$28,800 per QALY gained over 20 years by preventing diabetes-related complications and death. Compared with the EUC strategy, the CHW-only strategy was not cost-effective (\$430,600 per QALY gained) because of the dissipation of the intervention effects between 6 and 18 months. The

effectiveness of the PLs in maintaining the benefits of the intensive CHW-led DSME program, and its relatively low cost, accounted for it being cost-effective. When compared with the CHW-only strategy, the CHW + PL strategy is highly cost-effective (\$5,900 per QALY gained).

Our study is among the first to use a validated computer simulation model that is based on information from a short-term randomized controlled trial to evaluate the long-term cost-effectiveness of a CHW intervention. A systematic review (23) identified three published cost-effectiveness analyses of CHW interventions for T2D management using computer simulation modeling. They showed that a CHW intervention compared with usual care costs from \$371 to \$127,000 per QALY gained over a 20-year simulation time horizon from health system or societal perspectives. In contrast, the CHW-only intervention was not cost-effective in our base-case analysis. Given that two of the previously published studies mentioned above used very different designs, we only consider the third Community Diabetes Education (CoDE) (36) study for comparison. First, the CHW-led DSME intervention in the REACH Detroit trial was only 6 months, shorter than the 12-month intervention period in the CoDE trial. In the REACH Detroit trial, only brief CHW telephone contacts were made between 7 and 12 months followed by no further CHW contact between 13 and 18 months, which likely contributed to the waning effects in this group. Compared with the EUC intervention at 18 months, the CHW-only intervention was associated with lower HbA_{1c} and lipid levels but higher blood pressure levels, leading to a mixed and smaller impact on the incidence of some diabetes complications (e.g., stroke, congestive heart failure). This translated into a tiny improvement in QALYs, resulting in a large ICER for the CHW intervention compared with the EUC intervention in our base-case analysis.

In the base-case analyses, we found that relative to the EUC intervention, the CHW + PL intervention was cost-effective, but the CHW-only intervention was not. The improvements attributable to the 6-month CHW-led DSME intervention in risk factors, including HbA_{1c}

and SBP, were maintained or even greater in the CHW + PL group but not maintained at 18 months in the CHW-only group. This led to a lower incidence of diabetes-related complications in the CHW + PL group compared with the CHW-only group (Table 1), which averted downstream outcome costs and produced greater QALYs (Table 2). The additional support provided by PLs to the CHW + PL intervention group came at a relatively low cost. It follows that compared with the EUC intervention, the CHW + PL intervention was more economically attractive than the CHW-only intervention.

When we assumed that the benefits observed at 18 months after randomization persisted over 20 years, the CHW-only intervention was cost saving compared with the EUC intervention if no additional interventions were needed and cost-effective (\$22,700 per QALY gained) if limited additional interventions were needed. The assumption that the benefits observed at 18 months after randomization persisted over the 20-year simulation period is likely to be overly optimistic. The estimated cost-effectiveness of the CHW-only intervention relative to the EUC intervention was sensitive to assumptions regarding the persistence of the intervention benefits.

Some limitations to our study should be acknowledged. First, no simulation model can perfectly represent reality, and all models have inherent limitations (37). The validity of our modeling results are contingent on data quality and model assumptions. Misspecification of model parameters and structure are generally the most important sources of uncertainty. We have minimized first-order (stochastic) uncertainty and population uncertainty. To evaluate second-order (parameter) uncertainty, we also conducted sensitivity analyses to vary the costs and the treatment effects of the CHW + PL intervention. Our previous studies have demonstrated very good performance of the MMD (25). For this study, we used version 2 of the MMD, recalibrated the model parameters, and performed both internal and external validation studies to verify the model's performance. Second, the results that we simulated were based on a clinical trial and may not represent diabetes management in real-world

practice. Our study participants were Latino adults with T2D from a low-income urban setting. This may limit the generalizability of our findings. However, because Latino Americans experience a greater burden of diabetes complications than other racial/ethnic groups, the results obtained through this simulation study may have underestimated the long-term health and economic benefits (36). Finally, the cost-effectiveness analyses from the societal perspective were not performed because the cost data were unavailable from the informal health care and non-health care sectors, as shown in the impact inventory (Supplementary Table 4).

In summary, for Latino adults with T2D and poor glycemic control in a low-income urban setting, we found that adding a 12-month, less intensive PL-led DSME intervention after a 6-month CHW-led DSME intervention was cost-effective relative to both EUC and a CHW-only intervention. In some sensitivity analyses, this hybrid CHW + PL-led DSME/DSMS intervention could be cost saving. A 6-month intensive CHW-led DSME intervention with only limited follow-up was not cost-effective compared with EUC. Our pilot study of PL-led DSMS contributes to the literature supporting the use of such hybrid interventions. Larger clinical trials with longer follow-up are warranted to confirm and generalize these findings. From the perspective of a financially prudent policymaker, the CHW + PL intervention represents a good value for money. Health policy should support funding and reimbursement models for CHW + PL-led intervention programs.

Funding. The project described was supported by National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) grant P30-DK-092926 (Michigan Center for Diabetes Translational Research, Methods and Measurement Core), American Association of Family Physicians Foundation Peers for Progress grant, NIDDK grant R18-DK-0785501A1, and Centers for Disease Control and Prevention Cooperative Agreement No. U50/CCU417409.

Duality of Interest. No potential conflicts of interest relevant to this article were reported.

Author Contributions. W.Y., S.K., E.C.K., G.Pi., G.Pa., M.S.S., and W.H.H. contributed to the study concept and design. W.Y., S.K., E.C.K., and G.Pi. contributed to the acquisition, analysis, or interpretation of data. W.Y., S.K., and B.S. contributed to the statistical analysis. W.Y., E.C.K., and W.H.H. provided

study supervision. E.C.K. and W.H.H. contributed to the critical revision of the manuscript for important intellectual content. All authors contributed administrative, technical, and material support and to the drafting of the manuscript. W.Y., S.K., and W.H.H. are the guarantors of this work and, as such, had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Prior Presentation. Parts of this study were presented at the 78th Scientific Sessions of the American Diabetes Association, Orlando, FL, 22–26 June 2018.

References

- Centers for Disease Control and Prevention. National Diabetes Statistics Report, 2017. Atlanta, GA, Centers for Disease Control and Prevention, U.S. Department of Health and Human Services, 2017
- American Diabetes Association. Economic costs of diabetes in the U.S. in 2017. *Diabetes Care* 2018;41:917–928.
- Norris SL, Lau J, Smith SJ, Schmid CH, Engelgau MM. Self-management education for adults with type 2 diabetes: a meta-analysis of the effect on glycemic control. *Diabetes Care* 2002;25:1159–1171.
- Polonsky WH, Earles J, Smith S, et al. Integrating medical management with diabetes self-management training: a randomized control trial of the Diabetes Outpatient Intensive Treatment program. *Diabetes Care* 2003;26:3048–3053.
- Spencer MS, Rosland AM, Kieffer EC, et al. Effectiveness of a community health worker intervention among African American and Latino adults with type 2 diabetes: a randomized controlled trial. *Am J Public Health* 2011;101:2253–2260.
- Spencer MS, Kieffer EC, Sinco B, et al. Outcomes at 18 months from a community health worker and peer leader diabetes self-management program for Latino adults. *Diabetes Care* 2018;41:1414–1422.
- American Diabetes Association. (4) Foundations of care: education, nutrition, physical activity, smoking cessation, psychosocial care, and immunization. *Diabetes Care* 2015;38(Suppl.):S20–S30.
- Anderson G, Horvath J. The growing burden of chronic disease in America. *Public Health Rep* 2004;119:263–270.
- Eng E, Young R. Lay health advisors as community change agents. *Fam Community Health* 1992;15:24–40.
- Dennis CL. Peer support within a health care context: a concept analysis. *Int J Nurs Stud* 2003;40:321–332.
- Kieffer EC, Willis SK, Odoms-Young AM, et al. Reducing disparities in diabetes among African-American and Latino residents of Detroit: the essential role of community planning focus groups. *Ethn Dis* 2004;14(Suppl. 1):S27–S37.
- Feathers JT, Kieffer EC, Palmisano G, et al. The development, implementation, and process evaluation of the REACH Detroit partnership's diabetes lifestyle intervention. *Diabetes Educ* 2007;33:509–520.
- Palmas W, Findley SE, Mejia M, et al. Results of the northern Manhattan diabetes community outreach project: a randomized trial studying a community health worker intervention to improve diabetes care in Hispanic adults. *Diabetes Care* 2014;37:963–969.
- Shah M, Kaselitz E, Heisler M. The role of community health workers in diabetes: update on current literature. *Curr Diab Rep* 2013;13:163–171.
- Haas L, Maryniuk M, Beck J, et al.; 2012 Standards Revision Task Force. National standards for diabetes self-management education and support. *Diabetes Care* 2012;35:2393–2401.
- Norris SL, Chowdhury FM, Van Le K, et al. Effectiveness of community health workers in the care of persons with diabetes. *Diabet Med* 2006;23:544–556.
- Tang TS, Nwankwo R, Whiten Y, Oney C. Outcomes of a church-based diabetes prevention program delivered by peers: a feasibility study. *Diabetes Educ* 2014;40:223–230.
- Heisler M, Vijan S, Makki F, Piette JD. Diabetes control with reciprocal peer support versus nurse care management: a randomized trial. *Ann Intern Med* 2010;153:507–515.
- Heisler M. Different models to mobilize peer support to improve diabetes self-management and clinical outcomes: evidence, logistics, evaluation considerations and needs for future research [retracted in: *Fam Pract* 2012;29:497]. *Fam Pract* 2010;27(Suppl. 1):i23–i32.
- Heisler M. Overview of peer support models to improve diabetes self-management and clinical outcomes. *Diabetes Spectr* 2007;20:214–221.
- de Vries L, van der Heijden AA, van 't Riet E, et al. Peer support to decrease diabetes-related distress in patients with type 2 diabetes mellitus: design of a randomised controlled trial. *BMC Endocr Disord* 2014;14:21.
- Two Feathers J, Kieffer EC, Palmisano G, et al. Racial and Ethnic Approaches to Community Health (REACH) Detroit partnership: improving diabetes-related outcomes among African American and Latino adults. *Am J Public Health* 2005;95:1552–1560.
- Jacob V, Chattopadhyay SK, Hopkins DP, et al. Economics of community health workers for chronic disease: findings from community guide systematic reviews. *Am J Prev Med* 2019;56:e95–e106.
- Zhou H, Isaman DJM, Messinger S, et al. A computer simulation model of diabetes progression, quality of life, and cost. *Diabetes Care* 2005;28:2856–2863.
- Ye W, Brandle M, Brown MB, Herman WH. The Michigan model for coronary heart disease in type 2 diabetes: development and validation. *Diabetes Technol Ther* 2015;17:701–711.
- Brandle M, Zhou H, Smith BR, et al. The direct medical cost of type 2 diabetes. *Diabetes Care* 2003;26:2300–2304.
- Coffey JT, Brandle M, Zhou H, et al. Valuing health-related quality of life in diabetes. *Diabetes Care* 2002;25:2238–2243.
- University of Michigan, Michigan Center of Diabetes Translational Research (MCDTR) Disease Modeling Group. Michigan Model for Diabetes User Manual; 2015. Accessed 25 March 2021. Available from http://diabetesresearch.med.umich.edu/peripherals/DiseaseModel/MDRTC%20Diabetes%20Model/UserManual_Michigan_Model_for_Diabetes_ver2.pdf
- Sanders GD, Neumann PJ, Basu A, et al. Recommendations for conduct, methodological practices, and reporting of cost-effectiveness analyses: second panel on cost-effectiveness in health and medicine. *JAMA* 2016;316:1093–1103.
- Zhang P, Brown MB, Bilik D, Ackermann RT, Li R, Herman WH. Health utility scores for people with type 2 diabetes in U.S. managed care health plans: results from Translating Research Into Action for Diabetes (TRIAD). *Diabetes Care* 2012;35:2250–2256.
- Clarke PM, Gray AM, Briggs A, et al.; UK Prospective Diabetes Study (UKPDS) Group. A model to estimate the lifetime health outcomes of patients with type 2 diabetes: the United Kingdom Prospective Diabetes Study (UKPDS) Outcomes Model (UKPDS no. 68). *Diabetologia* 2004;47:1747–1759.
- Barhak J, Isaman DJM, Ye W, Lee D. Chronic disease modeling and simulation software. *J Biomed Inform* 2010;43:791–799.
- Braithwaite RS, Meltzer DO, King JT Jr., Leslie D, Roberts MS. What does the value of modern medicine say about the \$50,000 per quality-adjusted life-year decision rule? *Med Care* 2008;46:349–356.
- Laupacis A, Feeny D, Detsky AS, Tugwell PX. How attractive does a new technology have to be to warrant adoption and utilization? Tentative guidelines for using clinical and economic evaluations. *CMAJ* 1992;146:473–481.
- Neumann PJ, Cohen JT, Weinstein MC. Updating cost-effectiveness—the curious resilience of the \$50,000-per-QALY threshold. *N Engl J Med* 2014;371:796–797.
- Prezio EA, Pagán JA, Shuval K, Culica D. The Community Diabetes Education (CoDE) program: cost-effectiveness and health outcomes. *Am J Prev Med* 2014;47:771–779.
- Herman WH. Diabetes modeling. *Diabetes Care* 2003;26:3182–3183.